

# On the threshold of the SIS epidemic model with random perturbation

H. El Mahjour <sup>1</sup>,  
A. Akhrif<sup>1</sup>, A. Lahrouz<sup>2</sup>, A. Settati<sup>1</sup>

<sup>1</sup> Laboratory of Mathematics and Applications, Department of Mathematics, Faculty of Sciences and Techniques, B.P. 416-Tanger Principale, Tangier, Morocco

<sup>2</sup>Laboratory of Computer Sciences, Statistics and Quality, Department of Mathematics, Faculty of Sciences Dhar-Mehraz B.P. 1796-Atlas, Fez, Morocco

May 7, 2014

# Table of contents

## 1 Introduction

- Deterministic SIS model
- Stochastic SIS model

## 2 State of the art and main results

- Definition of a threshold
- Gray et al. results on  $\mathcal{R}_S$  (case  $\sigma^2 \leq \beta$ )
- Mains results of our work

# Introduction

Different models exist to study the progress of an epidemic in a population: SIR, SIRS, SIS etc.

- $S \rightarrow$  Susceptible
- $I \rightarrow$  Infected

**Example :** SIS models infections such as common flu

# Deterministic SIS model

$$\begin{cases} dS = (\mu - \mu S - \beta SI + \gamma I) dt, \\ dI = (-(\mu + \gamma)I + \beta SI) dt, \end{cases} \quad (1)$$

where :

- $\mu \rightarrow$  birth rate (coinciding with death rate)
- $\beta \rightarrow$  infection coefficient
- $\gamma \rightarrow$  recovery rate

Constant population  $\Rightarrow S + I = 1$

# Stochastic SIS model

$$\begin{cases} dS = (\mu - \mu S - \beta SI + \gamma I) dt, \\ dI = (-(\mu + \gamma)I + \beta SI) dt, \end{cases}$$

$$\beta \rightsquigarrow \beta + \sigma dB$$

$$\begin{cases} dS = (\mu - \mu S - \beta SI + \gamma I) dt - \sigma S I dB, \\ dI = (-(\mu + \gamma)I + \beta SI) dt + \sigma S I dB, \end{cases} \quad (2)$$

Recently (2011), Gray et al. considered the precedent stochastic model (2)

With initial conditions  $(S_0, I_0)$  in the set  $\Delta = \{x \in \mathbb{R}_+^2; x_1 + x_2 = 1\}$ .  
To solve the system 2 Gray et al. [2] proved that is sufficient to study the following EDS :

$$dI = (-(\mu + \gamma)I + \beta I(1 - I)) dt + \sigma I(1 - I)dB. \quad (3)$$

# Definition

$\mathcal{R}_S$  determines the dynamics of the SIS system.

It depends on the system parameters.

$$\mathcal{R}_S = f(\beta, \gamma, \mu, \dots)$$

- ① persistence
- ② extinction

In deterministic case  $\rightarrow \mathcal{R}_S = \frac{\beta}{\mu + \gamma}$

In stochastic case  $\rightarrow \mathcal{R}_S = \frac{\beta}{\mu + \gamma + \frac{1}{2}\sigma^2}$

# Gray et al. results on $\mathcal{R}_S$ (case $\sigma^2 \leq \beta$ )

- $\mathcal{R}_S < 1 \longrightarrow$  extinction
- $\mathcal{R}_S > 1 \longrightarrow$  persistence

$$\mathcal{R}_S = 1 \longrightarrow ?$$



# Main results of this communication

## Theorem

Let  $(S_0, I_0) \in \Delta$ . If  $\mathcal{R}_S < 1$  then for all  $\kappa$  such that

$$0 < \kappa < \frac{2\beta(1 - \mathcal{R}_S)}{\sigma^2 \mathcal{R}_S}, \quad (4)$$

the solution  $I(t)$  obeys

$$\mathbb{E}I^\kappa(t) \leq I_0^\kappa e^{-\Lambda t}, \quad \text{where } \Lambda = -\kappa \left[ \frac{\beta}{\mathcal{R}_S} (\mathcal{R}_S - 1) + \frac{\kappa}{2} \sigma^2 \right] > 0. \quad (5)$$

Then the disease-free equilibrium state  $E_0$  is  $\kappa$ -th moment exponentially stable.

## Theorem

Let  $(S_0, I_0) \in \Delta$ . If  $\mathcal{R}_S = 1$  then for all  $\eta > 0$  and  $\varepsilon > 0$ , we have

$$\lim_{I_0 \rightarrow 0} \mathbb{P} \left( \sup_{0 \leq t \leq \eta} I(t) > \varepsilon \right) = 0, \quad (6)$$


that is, the disease-free equilibrium state  $E_0$  is stable in probability.

## Theorem

*For any initial values  $(S_0, I_0) \in \Delta$ , if  $\mathcal{R}_S = 1$  then the solution of SDE (3) obeys*





$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t I_s ds = 0$$

# Simulation



**Figure 1 :** Computer simulation of a single path of  $I(t)$  for the SDE model (3) with initial condition  $I_0 = 0.4$  and its corresponding deterministic model for the parameters:  $\mu = 0.5$ ,  $\beta = 0.902$ ,  $\gamma = 0.4$ ,  $\sigma = 0.2$ , then  $\mathcal{R}_0 > 1$  and  $\mathcal{R}_S = 1$ .

# The bibliography

-  J. Zhou, H.W. Hethcote, Population size dependent incidence in models for diseases without immunity, J. Math. Biol. 32 (1994) 809-834.
-  A. Gray, D. Greenhalgh, L. Hu, X. Mao, J. Pan, A Stochastic Differential Equation SIS Epidemic Model, SIAM J. Appl. Math. 71 (2011) 876-902.
-  A. Gray, D. Greenhalgh, X. Mao., J. Pan, The SIS epidemic model with Markovian switching, J. Math. Anal. Appl. 394 (2012) 496-516.
-  A. Lahrouz, A. Settati. Asymptotic properties of switching diffusion epidemic model with varying population size. Applied Mathematics and Computation, 219 (2013) 11134-11148.

Thank you !